

Spheroid-like Cultures for Cell Expansion of Angiopoietin Receptor-1 (aka. Tie2) positive Cells from the human Intervertebral Disc

Xingshuo ZHANG¹, Julien GUERRERO¹, Sonja HäCKEL², Andreas Shaun CROFT¹, Selianne GRAF¹, Andrea OBERLI¹, Daisuke SAKAI³, Marianna TRYFONIDOU⁴, Lorin Michael BENNEKER², Benjamin GANTENBEIN²

¹Tissue Engineering for Orthopaedics & Mechanobiology (TOM), Department for BioMedical Research (DBMR), University of Bern, Bern, Switzerland

²Department of Orthopaedic Surgery & Traumatology, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

³Department for Orthopaedic Surgery, Tokai University School of Medicine, Isehara, Kanagawa, Japan

⁴Clinical Sciences of Companion Animals, Faculty of Veterinary Medicine, Utrecht University, Utrecht, the Netherlands

INTRODUCTION:Low back pain is the leading cause of disability worldwide (1). Nevertheless, the mechanism of the intervertebral disc (IVD) degeneration is still not clear. In this context, the nucleus pulposus (NP) and more precisely NP progenitor cells (NPPCs) present in the IVD, positive for angiopoietin-1 receptor (aka. Tie2) display multipotent and stem capacity (2,3). In this study, the first aim was to determine whether spheroid formation in suspension-culture will increase the amount/percentage of NPPCs during the expansion compared to traditional monolayer culture. The second aim of this study was to investigate if the percentage of NPPCs will be enriched even further by the resuspension of the spheroid-like cultured cells (=1st generation) and reformation of those spheroids one more time (= 2nd generation).

METHODS:Human NP tissues from trauma patients (N=3) were obtained with written ethical consent and isolated by a two-step digestion protocol (3). The NP cells were resuspended and frozen at -150°C after reaching confluence of passage 0. At passage 1, NP cells were seeded in standard or ultra-low attachment tissue culture flasks with 2.5 ng/ml FGF-2 in low glucose - DMEM (supplemented with 10 % FBS). Flow cytometry was used to analyze and quantify the percentage of NPPCs using Tie2 antibody. We defined the spheroids formed after passage 1 NPCs as 1st generation spheroid. We obtained the 2nd generation spheroids by resuspending the 1st-generation-spheroid and reassembly. The NPCs from 1st and 2nd spheroid were quantified by CFU-assay.

RESULTS:As a result, the percentage of NPPCs in monolayer culture condition was reaching 7 ± 2 % (Mean \pm SEM), however, in the 1st and 2nd generation spheroids culture condition, we were observing 20 ± 10 % and 28 ± 6 % of Tie2+ cells, respectively. Concerning the CFU-assay, the NPCs from the 2nd generation spheroid formed 30 CFU-S per 1,000 cells, which were twice more CFU-S compared to the 1st generation spheroid.

DISCUSSION & CONCLUSIONS:From these data we conclude that the spheroid-like formation of NPCs would be a more efficient method for expansion and enrichment of NPPCs than monolayer expansion in a context of future cell therapy.

Acknowledgements:Financial support was received from iSpine H2020 project under grant agreement #825925 and China Scholarship Council to X.Z.

References:[1] Lancaster center. (2013)

[2] Sakai D et al. (2012)

[3] Frauchiger, D. A et al. (2019)

Keywords: Intervertebral disc / spine and their disorders, Stem cells – general